

Kaposi's sarcoma of a lymph node with Castleman's disease-like lesions in HIV-infected patient: A case report

Thamathorn Assanasen*

Pongsak Wannakrairot*

Assanasen T, Wannakrairot P. Kaposi's sarcoma of a lymph node with Castleman's disease-like lesions in HIV-infected patient: A case report. Chula Med J 2002 May; 46(5): 409 - 17

The histomorphology of Castleman's disease of lymph nodes is not specific only for Castleman's disease. Kaposi's sarcomas (KS) and HIV infection of lymph nodes are also found in a number of diseases. We, herein, report the rare characteristics found in the same lymph node of an HIV-infected patient, to remind pathologists about the existence of this histomorphology. The case report was of a Thai patient, admitted at King Chulalongkorn Memorial Hospital, who presented with multiple cervical lymphadenopathy. A lymph node biopsy and a routine histopathological examination were done. A coexistence, a Castleman's disease-like lesion and Kaposi's sarcoma, was observed in the same lymph node. The backgrounds of both lesions were reviewed with a discussion about their association and the roles of Human herpes virus 8 (HHV-8). According to the findings and reviewed papers, one should not diagnoses Castleman's disease without a prior extensive exclusion of other diseases.

Key words: Kaposi's sarcoma, Castleman's disease, Lymph node, HIV, HHV-8.

Reprint request : Assanasen T, Department of Pathology, Faculty of Medicine,
Chulalongkorn University, Bangkok 10330, Thailand.

Received for publication. February 12, 2002.

ธรรมธร อาศนะเสน, พงษ์ศักดิ์ วรรณไกรโรจน์. มะเร็งต่อมน้ำเหลืองชนิด Kaposi's sarcoma ที่มีลักษณะรูปร่างทางกล้องจุลทรรศน์ ของต่อมน้ำเหลืองคล้ายโรค Castleman ในผู้ป่วยติดเชื้อ HIV: รายงานผู้ป่วย 1 ราย. จุฬาลงกรณ์เวชสาร 2545 พ.ศ; 46(5): 409 - 17

ลักษณะทางกล้องจุลทรรศน์ ของต่อมน้ำเหลืองในโรค Castleman นั้นไม่ได้จำเพาะเฉพาะโรคนี้ อาจพบได้ในโรคอื่น ๆ รวมทั้ง Kaposi's sarcoma ของต่อมน้ำเหลือง และผู้ป่วยที่ติดเชื้อ HIV การรายงานผู้ป่วยรายนี้ มีเจตนาจะให้พยาธิแพทย์ระลึกถึงความจริงดังกล่าว ผู้ป่วยรายนี้ เป็นคนไข้ในโรงพยาบาลจุฬาลงกรณ์ ที่มาด้วยต่อมน้ำเหลืองที่คอโตหลายต่อม และได้ถูกตัดออกแล้วนำไปตรวจทางพยาธิวิทยา การรายงานนี้ยังรวมถึงการศึกษาลักษณะของทั้ง 2 โรคโดยละเอียด พร้อมกับความสัมพันธ์ซึ่งกันและกัน และความเกี่ยวข้องของ Herpes simplex virus 8 จากการศึกษาการจะวินิจฉัยถึงโรค Castleman นั้นต้องทำการแยกโรคอื่น ๆ ออกให้หมดก่อนจึงจะวินิจฉัย มิใช่ดูจากลักษณะทางกล้องจุลทรรศน์แต่เพียงอย่างเดียว

Kaposi's sarcoma (KS) was relatively rare in Asian countries, until the outbreak of AIDS epidemic. It has become the most common AIDS-related cancer reported internationally.⁽¹⁾

The pandemic of Kaposi's sarcoma of lymphadenopathic type among homosexual men⁽²⁾ and those who acquired immunodeficiency syndrome (AIDS),⁽³⁾ has pushed this, once a rare disorder among developed countries, (it was a common tumor in several parts of Africa) into the forefront of global medical interest.

We hereby report a case of KS with Castleman's disease-like features of a lymph node in Human Immunodeficiency Virus (HIV), taken from an infected patient (at King Chulalongkorn Memorial Hospital). Our purpose is to illustrate histopathological features of Castleman's disease-like lesion and KS, which coexisted in a lymph node and the difficulties in establishing the diagnosis.

Case report

A 34-year-old Thai male, presented at King Chulalongkorn Memorial Hospital in May 2000, with multiple cervical lymphadenopathy hepatosplenomegaly and skin rash without fever. His serum anti-HIV was positive. A cervical lymph node biopsy was taken and sent to Department of Pathology, Faculty of Medicine, Chulalongkorn University for histological examination. The macroscopic appearance of the lymph node was as follows: a well-circumscribed firm ovoid brown mass, measured 1.5x1x0.5 cm. Its cut surfaces were pinkish brown with multiple red brown spots. Its microscopic examination revealed a preserved nodal architecture with scattered multiple discrete angiomatous lesions and slit-like vascular

spaces; extravasated erythrocytes and an intervening spindle cells' components were found in the lymph node with some areas adjacent to its capsule. Involvement of sinuses was seen with focal parenchymal replacement. The spindle cells possessed elongated nuclei with pale to dark chromatin, inconspicuous nucleoli, and lightly stained cytoplasm. Nuclear pleomorphism was mild, and mitosis was rarely seen (Fig.1). There were some hyaline globules (2 - 10 μ m) that were PAS-positive and diastase-resistant. Most of them were located within the cytoplasm of the spindle cells or histiocytes, surrounded by clear halos. Occasionally, the globules were extra-cellular. The residual lymphoid tissue displayed abnormal follicles and interfollicular vascularity. The follicles were increased, and varied in sizes, from barely perceptible




Figure 1. Kaposi's sarcoma in lymph node: A high magnification shows typical criss-crossing pattern, with longitudinal sectioned fascicles alternated with transversely sectioned fascicles. Note sieve-like pattern in the latter. Nuclear pleomorphism is minimal and red blood cells are seen in vascular slits (H&E, x 200).

to medium, but most of them were small. Many follicles contained small blood vessels that often radiated from the perifollicular vessels through the germinal centers. The capillaries had thick hyalinized walls, which gave rise to thickened hyaline in the follicles. Other follicles were characterized by concentric layering of cells within germinal centers, thus superficially resembling Hassle's corpuscles of the thymus. Some of the follicular centers were surrounded by concentric zones ("onion skin"), small lymphocytes (Fig. 2) which were prominent with completely obscured germinal centers and presence of more than one small germinal center within a single

follicle in few follicles. The inter-follicular areas consisted of extensive networks of capillaries. The majority of the cells among these capillaries were plasma cells (Fig.3), with occasional immunoblasts. Few small fibrotic areas were also noted. The acid-fast stain showed no positive acid-fast bacilli. The neoplastic spindle cells showed focally positive staining for factor VIII, actin and CD34. The final anatomical diagnosis of the lymph node was Kaposi's sarcoma of the lymph node with Castleman's disease-like features. The study for HHV-8 was not available at our institute.




Figure 2. Hyaline-vascular type of Castleman's like features: There is concentric layering of small lymphocytes around a compact germinal center, devoid of follicle center cells, which superficially resembling a Hassal's corpuscle. A capillary from the vascular interfollicular area penetrates the follicle (H&E, x100).




Figure 3. The interfollicular area contains an almost pure population of plasma cells (H&E, x 400).

Discussion

Kaposi's sarcoma (KS) is a distinctive tumor of probable vascular nature. Histologically, it is characterized by an admixture of spindle cells and narrow vascular spaces, and frequently showing multicentric growths.⁽⁴⁾ In the case, there was no involvement of other organ, except the lymph node. It was controversy concerning the nature of KS, whether it was a hyperplastic lesion or a neoplasm.⁽⁵⁾ The multifocal growth, which did not conform to a metaplastic pattern, the heterogeneous cellular composition, and the lack of DNA aneuploidy were cited to support the probable hyperplastic nature of KS.⁽⁶⁾ On the other hand, in vitro studies demonstrated transforming genes in KS.⁽⁷⁾ DNA obtained from NIH/3T3, when the transfected cells were injected into nude mice; a tumor that was histologically similar to KS was formed. It was possible that early KS was a hyperplastic state that might progress into a true neoplasm, when a secondary genetic change occurred.⁽⁸⁾ Kaposi's sarcoma occurred in a variety of clinical settings, but it was rare among the non-Africans until the recent epidemic of AIDS.⁽⁹⁾ KS occurred in approximately 30 percent of patients with AIDS, primarily among homosexual men; however, it was uncommon among patients who contracted HIV infection through parenteral route.⁽¹⁰⁾ The risk of developing KS was fairly constant after HIV seroconversion, unlike the risk of opportunistic infections or lymphoma which increased as the immunodeficiency became worsen. The disorder was often disseminated, involving mucocutaneous sites, lymph nodes, and visceral organs (especially the gastrointestinal tract and lungs). AIDS-related KS also had a fulminant course, with less than 20 percent

survival in 2 years, if it had opportunistic infection.⁽¹¹⁾

A recent study revealed human herpes virus 8 (HHV-8) in early lesions supporting a role for HHV-8 in the pathogenesis of KS.⁽¹²⁾ Furthermore, detection of HHV-8 in HIV-positive patients without KS was often predictive of a subsequent development of KS. A serologic study showed that 52% of patients with KS had seroconversion of antibody to HHV-8, with the median time of 33 months before the diagnosis of KS. The mechanism of oncogenesis of HHV-8 might be related to the ability of HHV-8 to encode homology of Bcl-2 chemokine receptor, interleukin-6, and cycline D.

Castleman's disease was first described by Castleman and associates.⁽¹³⁾ It had many synonyms,⁽¹⁴⁾ namely: = lymph nodal hamartoma, angiofollicular mediastinal lymph node hyperplasia, angiomatous lymphoid hamartoma, follicular lymphoreticuloma, and benign giant lymphoma. In the first accounts of giant lymph node hyperplasia, the lesions were described as solitary and localized to the mediastinum, which was still the most frequent site of its involvement. Giant lymph node hyperplasia was reported as a solitary lesion in other locations, .eg., cervical and axillary regions, skeletal muscle, pulmonary parenchyma, abdomen, and retroperitoneum.⁽¹⁵⁾ There appeared to be no sex preference, and the age of patients ranged from 8 to 70 years. Two distinct clinical manifestations were recognized. Solitary Castleman's disease and Multicentric Castleman's disease (MCD). The former seemed to be non-specific; and its symptoms were usually due to its mass effects. It usually consisted of a rounded mass ranging in size from 1.5 - 16.0 cm⁽¹⁷⁾ MCD, on the other hand, was a systemic lymphoproliferative disorder with morphologic features of Castleman's

disease. The diagnosis of this disease came out of exclusion of other disorders that had similar histological patterns. Its patients were older (57 years, median age) than those with the localized disease, most of them were male, and their peripheral lymph nodes were always involved, and the development of malignancy was common, and its clinical course of the disorder varied. It might behave like a chronic disease with persistent or relapsing pattern, otherwise like an unremitting aggressive lymphoproliferative disorder that might be fatal. The disorder might associate with KS, POEMS, or Crow Fukase syndrome. Lymph nodes with similar histologic changes were also reported in patients with AIDS. Other organs frequently involved by systemic disorder included bone marrow, liver, kidney, skin, spleen, and central and peripheral nervous system.⁽¹⁶⁾ Histologically, there were two distinct types of lesions described: hyaline-vascular and plasma cell. A transitional or an intermediate form of pathology with similar features of both was also reported.⁽¹⁶⁾ The hyaline-vascular type of lesions was much more common than the plasma cell type. The majority of patients with the hyaline-vascular type of lesion were asymptomatic except for mass effects. In contrast, patients with plasma cell type of giant lymph node hyperplasia, which constituted about 10 -20 percent of the lesions, often had associated systemic clinical manifestations, such as fever, anemia, and hypergammaglobulinemia.⁽¹⁷⁾ The symptoms were reported to disappear after the removal of the lesions. The transitional or mixed type of Castleman's disease was to be the least common.⁽¹⁴⁾ Clinically, the majority of patients were similar with those who had hyaline vascular type in that they were asymptomatic at their presentation.

Histologically, most of the lesions were of the hyaline-vascular type with focal areas of plasmacytosis. The etiology of the lesions was not known. The changes in hyaline vascular type suggested an inflammatory or infectious process, perhaps a later stage of the plasma cell type.⁽¹⁴⁾ Excessive interleukin-6 (IL-6)-like activity was found. The data suggested that Castleman's disease was a syndrome of multiple etiologies with aberrant IL-6 activity from either endogenous or viral sources.⁽¹⁸⁾ HHV-8 was also associated with MCD. HHV-8 was found in most cases of HIV-positive MCD. Even in HIV-negative patients HHV-8 appeared to have significant association with MCD, though the findings were not striking. The role of HHV-8 in pathogenesis of MCD remained unclear.⁽¹⁹⁾

HHV-8 played a striking role in the pathogenesis of both KS and MCD. HHV-8 was first isolated by Chang et al, 1994. The unique DNA sequences were isolated from KS tissue from AIDS patients. The KS-associated herpes-virus like sequences (KSHV) had significant homology to sequences of two gamma herpes viruses; EBV and herpes virus saimiri, then was named Human herpes virus 8 (HHV-8). In the subsequent studies, HHV-8 appeared to be associated with all types of KS, whether the patients were antiHIV-positive or negative, or the patients were antiHIV-positive via homosexual transmission, or other routes.⁽²⁰⁻²²⁾

The association between MCD and KS was studied. In antiHIV-negative patients with MCD, there was 13 % chance of KS during the course of the disease, but in antiHIV-positive patients with MCD, 75 % developed KS.⁽²³⁾ It was unclear whether KS was a consequence of immunological alterations associated with MCD, or whether both processes

were caused by a primary disorder of immune regulation.⁽²⁴⁾ In the case, only one lymph node was examined. Multicentricity was not verified as though investigation of the whole body were not carried out. The study of HHV-8 was not available at our institute.

It is worth to mention that features that characterize Castleman's disease, especially hyaline vascular type, which could be found in other conditions, such as AIDS-related lymphadenopathy, systemic lupus erythematosus, angioimmunoblastic lymphadenopathy, and single follicle, otherwise in non-specific reactive lymph nodes.^(8,14,25,26) There was an example of uninvolved areas of lymph nodes in KS which showed features of giant lymph node hyperplasia of Castleman (hyaline-vascular type).^(27,28) The term Castleman's disease-like lesion might be applied to a node without any obvious clinical manifestation of Castleman's disease.

References

1. Bhoopat SK, Vithayasai P, Prutasachatkenakorn C. AIDS-- associated Kaposi's sarcoma: a rare entity at Maharaj Nakorn Chiang Mai Hospital. *Asian Pac J Allergy Immunol* 1996 Dec; 14(2): 115 - 20
2. Finkheiner WE, Egberst BM, Groundwater JR, Sagabiel RW. Kaposi's sarcoma in young homosexual men: a histologic study with particular reference of lymph node involvement. *Arch Pathol Lab Med* 1982 Jun; 106(6): 261 - 4
3. Safai B, Johnson KG, Myskowski PL, Koziner B, Vang SY, Cunningham-Rundles S, Godbold JH, Dupont B. The natural history of Kaposi's sarcoma with the acquired immunodeficiency syndrome. *Ann Intern Med* 1985 Nov; 103(5): 744 - 50
4. Templeton AC. Pathology of Kaposi's sarcoma. In: Ziegler JL, Dorfman FR, eds. *Kaposi's Sarcoma, Pathophysiology and Clinical Management*. New York: Marcel Dekkes, 1988: 23 - 70
5. Costa J, Rabson AS. Generalized Kaposi's sarcoma is not a neoplasm [letter]. *Lancet* 1983 Jan 1; 1(8314-5): 58
6. Auerbach HE, Brooks JJ. Kaposi's sarcoma: neoplasia or hyperplasia? *Surg Pathol* 1989; 2 :19 - 28
7. Lo SC, Liotta LA. Vascular tumours produced by NIH/3T3 cells transfected with human AIDS Kaposi's sarcoma DNA. *Am J Pathol* 1985 Jan; 118(1): 7 - 13
8. Ensoli B, Salaholdin SZ, Gallo RC. AIDS-associated Kaposi is sarcoma. A molecular model for its pathogenesis. *Cancer Cells* 1989 Nov; 1(3): 93 - 96
9. Gottlieb GJ, Ackerman AB. Kaposi's sarcoma: an extensively disseminated form in young homosexual men. *Hum Pathol* 1982 Oct; 13(3): 882 - 92
10. Beral V, Peterman TA, Berkeleman RL, Jaffe HW. Kaposi's sarcoma among persons with AIDS: a sexually transmitted infection? *Lancet* 1990 Jan 20; 335(8682): 123 - 8
11. Warnke RA, Wiess LM, Chan JKC. Nonhematolymphoid tumors and tumor-like lesions of lymph node and spleen. In: Rosia J, ed. *Atlas of Tumor Pathology "Tumours of the Lymph nodes and Spleen"*. Washington D.C. Armed Forces Institute of Pathology, 1995: 435 - 526

12. Kennedy MM, Cooper K, Howells DD, Picton S, Biddolps S, Lucas SB, McGeeJo, O'Leary JJ. Identification of HHV8 in early Kaposi's sarcoma: implications for Kaposi's sarcoma pathogenesis. *Mol Pathology* 1998 Feb; 51 (1): 14 - 20
13. Castleman B, Iverson L, Menendez VP. Localized mediastinal lymph node hyperplasia resembling thymoma. *Cancer* 1956 Jul - Aug; 9(4): 822 - 30
14. Schnitzen B. Reactive lymphoid hyperplasia. In: Jeffe ES, ed. *Surgical Pathology of the Lymph Nodes and Related Organs*. 2nd. Philadelphia: W.B. Saunders, 1995: 98 - 132
15. Tung KSK, Mc Cormack LJ. Angiomatous lymphoid hamartoma. Report of five cases with a review of the literature. *Cancer* 1967; 20: 525 - 36
16. Gaba AR, Stein RS, Sweet DL, Varialcojis P. Multicentric giant lymph node hyperplasia. *Am J Clin Pathol* 1978 Jan; 69(1): 86 - 90
17. Keller AR, Hochholzer L, Castleman B. Hyaline-vascular and plasma-cell types of giant lymph node hyperplasia of mediastinum and other locations. *Cancer* 1972 Mar; 29(3): 670 - 83
18. Parravinci C, Corbellino M, Paulli M, Magrini U, Lazzarino M, Moore PS, Chang Y. Expression of a virus-derived cytokine, KSHV VIL-6, in HIV-seronegative Castleman's disease. *Am J Pathol* 1997 Dec; 151(6): 1517-22
19. Corbellinu M, Poirel L, Aubin JT, Paulli M, Magrini U, Bestetti G, Galli M, Panavicini G. The role of human herpesvirus 8 and Epstein-Barr virus in the pathogenesis of giant lymph node hyperplasia (Castleman's disease). *Clin Infect Dis*. 1996 Jun; 22(6): 1120 - 1
20. Dupin N, Fisher C, Kellam P, Ariad S, Tulliez M, Franck N, van Mark E, Salmon D, Gorin I, Escande JP. Distribution of human Herpesvirus-8 latently infected cells in Kaposi's sarcoma, multicentric Castleman's disease, and primary effusion lymphoma. *Proc Natl Acad Sci USA*. 1999 Apr 13; 96(8): 4546 - 51
21. Oksenhendler E, Crazals-Hatem O, Schulz TR, Barateau V, Grollet L, Sheldon J, Glauvel JP, Sigaux F, Agbalika F. Transient angiolymphoid hyperplasia and Kaposi's sarcoma after primary infection with human herpesvirus 8 in a patient with human immunodeficiency virus infection. *N Engl J Med*. 1998 May 28; 338(22): 1585 - 90
22. Nampoory MR, Johnny KV, Sarkar C, Al-Masry I, Al-Hilali N, Anian JT. The dialysed patient with both Castleman's disease and Kaposi's sarcoma: *Nephrol Dial, Transplant* 1998 Sep; 13(9): 2373 - 6
23. Oksenhendler E, Duarte M, Soulier J, Cacoub P, Welker Y, Gadranel J, Gazals-Hatem D, Autran B, Glauvel JP, Raphael M. Multicentric Castleman's disease in HIV infection: a clinical and pathological study of 20 patients. *AIDS* 1996 Jan; 10(1): 61 - 7
24. Chen KT. Multicentric Castleman's disease and Kaposi's sarcoma. *Am J Surg Pathol* 1984 Apr; 8(4): 287 - 93
25. Lowenthal OA, Fillipa DA, Richardson ME, Bertoni M, Straus DJ. Generalized lymphadenopathy with morphologic features of Castleman's disease in an HIV-positive man. *Cancer* 1987 Nov 15; 60(10): 2454 - 8
26. Sohn CC, Sheibani K, Winberg CA, Rappaport H.

Monocytoid B lymphocytes: their relation to the patterns of the acquired immunodeficiency syndrome (AIDS) and AIDS-related lymphadenopathy. Hum Pathol 1985 Oct; 16(10): 979 - 85

27. Harris NL. Hypervascular follicular hyperplasia and Kaposi's sarcoma in patient at risk for AIDS. N Engl J Med 1984 Feb 16; 310(7): 462 - 3

28. Perlow LS, Paff ML, Orsini JM, Goldsmith MA, Hruza ZT, Gerber MA, Geller SA. Kaposi's sarcoma in a young homosexual men. Associated with angiofollicular lymphoid hyperplasia and a malignant lymphoproliferative disorder. Arch Pathol Lab Med 1983 Oct; 107(10): 510 - 3